

SPOT DIAGNOSIS (IMAGE GALLERY)



MULTIPLE SOFT TISSUE CALCIFICATIONS WITH TOE DEFORMITIES

Chandra Kanta, Rashmi Kumar

Department of Pediatrics, KG Medical University, Lucknow, Uttar Pradesh, India.

Address for Correspondence: Dr. Chandra Kanta, Department of Pediatrics, KG Medical University, Lucknow, Uttar Pradesh, India

A ten years old female child, born of consanguineous marriage, presented with bilateral toes deformities since birth and multiple bony swellings over neck and trunk increasing progressively in number and size for three years.

What is the diagnosis?

Fibrodysplasia ossificans progressiva (FOP) is a rare disabling hereditary disorder of connective tissues characterized by symmetric congenital anomalies of the great toes and thumbs with progressive post-natal heterotopic ossification of soft tissues including those related to the striated muscles, leading to permanent disabilities. (1) It affects men and women equally, with a current worldwide prevalence of approximately 1 case in 2 million individuals. (2) Calcification of soft tissues starts during the first decade of life at 5 years on average. (3) Episodes of painful soft tissue swellings occur which are often precipitated by soft tissue injury, intramuscular injections, viral infection, muscular stretching, falls or fatigue. These flare-ups transform skeletal muscles, tendons, ligaments, fascia, and aponeuroses into heterotopic bone, rendering movement impossible. (3) Although most cases of FOP are sporadic (non-inherited mutations), a small number of inherited FOP cases show germline transmission in an autosomal dominant pattern. Heterozygous activating mutations in activin receptor IAPerctivin-like kinase-2 (ACVRI, ALK2), a bone morphogenetic protein (BMP) type I receptor, exist in all sporadic and familial cases of FOP. (1,2) Differential diagnosis includes progressive osseous heteroplasia, osteosarcoma, lymphedema, soft tissue sarcoma, desmoid tumors, aggressive juvenile fibromatosis, and non-hereditary (acquired) heterotopic ossification. Skeletal survey is required to find out extent of involvement. (4) Current management is focused on early diagnosis, assiduous avoidance of injury or iatrogenic harm, symptomatic amelioration of painful flare-ups, and optimization of residual function. Short course steroid is helpful during acute exacerbations. Treatment with bisphosphonate and isotretinoin help to some extent but disease is slowly progressive and most of the patients are bed ridden by 30 years. (5,6) Death occurs by second to fourth decade usually due to respiratory failure. (3)

References

1. Feldman G, Li M, Martin S, Urbanek M, Urtizberea JA, Fardeau M, et al. Fibrodysplasia ossificans progressiva: A heritable disorder of severe heterotopic ossification, maps to human chromosome 4q27-31. *Am J Hum Genet.* 2000;66:128-135
2. Shore EM, Xu M, Feldman GJ, Fenstermacher DA, Cho TJ, Choi IH, et al. A recurrent mutation in the BMP type I receptor ACVR1 causes inherited and sporadic fibrodysplasia ossificans progressiva. *Nature Genetics* 2006; 38:525-527
3. Cohen RB, Hahn GV, Tabas JA, Peeper J, Levitz CL, Sanda A, et al. The natural history of heterotopic ossification in patients who have fibrodysplasia ossificans progressiva: A study of forty-four patients. *J Bone Joint Surg Am.* 1993;75:215-219
4. Cremin B, Connor MJ, Beighton P. The radiological spectrum of fibrodysplasia ossificans progressiva. *Clin Radiol.* 1982;33:499-508
5. Kaplan FS, Glaser DL, Pignolo RJ, Shore EM. A new era for fibrodysplasia ossificans progressiva: A druggable target for the second skeleton. *Exp Opin Biol Ther.* 2007; 7: 705-712.
6. Illingworth RS. Myositis ossificans progressiva: Brief review with report of two cases treated with corticosteroids and observed for 6 years. *Arch Dis Child.* 1971;46:264-268

DOI No. : 10.7199/ped.oncall.2014.20

Quick Response Code

