
SPOT DIAGNOSIS (IMAGE GALLERY)



SKELETAL ANOMALY

Vikrant Sood, Mukul Aggarwal ,

Department of Pediatrics, Vardhman Mahavir Medical College and Safdarjang Hospital, New Delhi, India

Address for Correspondence: Dr Vikrant Sood, Vardhman Mahavir Medical College and Safdarjang Hospital, New Delhi, India. Email: drvickyster@gmail.com

Absent radius. It is commonly seen with Thrombocytopenia-absent radius {TAR}, Fanconi Syndrome, Holt-Oram Syndrome, Rapadilino Syndrome, Roberts Syndrome, Thalidomide Embryopathy, Trisomy 18 {Edward Syndrome}, And Vacterl Association. Absent radii is detectable during prenatal transvaginal ultrasonography as early as 13 weeks` gestation. Treatment includes splinting of the hands during infancy. Adaptive devices to assist with activities of daily living {e.g. dressing, toileting, feeding} are helpful. Prostheses are less useful than adaptive devices.

References:

1. Caumes E, Ly F, Bricaire F. Cutaneous larva migrans with folliculitis: report of seven cases and review of the literature. Br J Dermatol. 2002; 146: 314 -316
2. Blackwell V, Vega-Lopez F. Cutaneous larva migrans: clinical features and management of 44 cases presenting in the returning traveler. Br J Dermatol. 2001; 145: 434 -437

E-published: January 2011 . **Art#5**