

TEACHING FILES (GRAND ROUNDS)

IS IT GAUCHER'S DISEASE OR NIEMANN-PICK'S DISEASE?

Lavina Desai¹, Ira Shah²

¹Seth G S Medical College, Mumbai, India and ²Department of Pediatric Hepatology, Nanavati Hospital, Mumbai, India.

KEYWORDS

Gaucher, NPC, GD

ARTICLE HISTORY

Received 9 June 2018

Accepted 23 October 2018

Clinical Problem

A three and a half-year-old girl, currently bedridden, presented in 2013 with progressive loss of milestones from two years of age. On examination she showed poor oral hygiene, oral thrush over the hard palate, left cervical lymphadenopathy of size 2x2cm, and hepatomegaly (1cm), splenomegaly (10 cm). In 2011, she was diagnosed with Gaucher's disease and advised regarding enzyme replacement therapy. Bone marrow examination at that time had shown foamy macrophages and a lysosomal enzyme study revealed reduced levels of beta-glucosidase (Patient: 3.7 nmol/m/hr, Control: >13 nmol/m/hr), raised plasma chitotriosidase (4274.8 nmol/m/hr) and normal levels of sphingomyelinase enzyme. She also had pancytopenia. In July 2012, a genetic test for Gaucher's disease was negative. On presentation to us in 2013 in view of neurological involvement, a repeat beta-glucosidase levels were done which was 3 nmol/m/hr (low), chitotriosidase was 2879 nmol/m/hr and sphingomyelinase was 9.3 nmol/m/hr (normal). A further genetic test showed NPC1 gene mutation and a normal NPC2 gene suggestive of Niemann-pick type C 1 disease.

Is this Gaucher's disease or Niemann Pick disease?

Discussion

Gaucher's disease and Niemann-Pick's (NPC) disease are classified as lysosomal storage disorders with autosomal recessive inheritance. Gaucher's disease (GD) is the most common lysosomal storage disease in India and worldwide and should be considered in any child or adult with unexplained hepatosplenomegaly and cytopenia. (1) It is caused due to deficiency of beta-glucosidase and leads to accumulation of lipid-laden macrophages in various tissues. Type 1 GD is a non-neuronopathic form and type 2 and 3 are neuronopathic (1). Niemann pick disease (NPD) is classified as Type A (Classic infantile neuronopathic form), Type B (Non-neuronopathic visceral) and Type C (juvenile form). (2) Type A and Type B occur due to deficiency of acid sphingomyelinase. Type C occurs due

to defective transport of cholesterol and glycolipid. (3) They ultimately lead to accumulation of sphingomyelin and cholesterol in the monocyte-macrophage system characterized by lipid-laden macrophages or Niemann pick cells (NP cells). (2) Niemann-pick disease type C (NPC) in approximately 95% of patients have a mutation in NPC1 gene which encodes large membrane glycoproteins, and the remainder 5% have mutations in NPC2 gene which encodes a small soluble lysosomal protein with cholesterol binding properties. (4)

Type 1 GD and Type1 NPD have overlapping presentation. They usually present from infancy to late childhood with hepatosplenomegaly, cytopenias, irritability, bone marrow involvement and failure to thrive. NPD additionally also causes loss of early motor skills which was seen in our patient, recurrent infections, and in later stages, loss of intellectual capabilities, rigidity and spasticity. (5,6) Definitive diagnosis for both diseases is made by enzyme assay. Molecular studies for mutation are useful in confirming diagnosis, screening family members, and prognosticating the disease. (1)

Our patient was initially diagnosed with GD due to reduced levels of beta-glucosidase, elevated chitotriosidase, normal levels of sphingomyelinase, and compatible clinical presentation like hepatosplenomegaly and pancytopenia. Low beta-glucosidase activity is observed in both NPD and GD, at least in leukocytes and can lead to false positive testing for GD. Glucocerebrosidase (GCase) accumulates within cells deficient in NPC1 protein and this occurs despite the lack of GBA1 mutation. The precursor of glucocerebrosidase/beta-glucosidase is found at normal levels in NPC fibroblasts, however, the level of the mature protein is reduced. This pattern suggests accelerated breakdown of mature GCase rather than decrease in production. This is in contrast to GD1 where the GBA1 gene mutation causes a decrease in production of GCase. (7) A second diagnostic marker like chitotriosidase or TRAP can help in diagnosis. (7) Chitotriosidase (CT) is an enzyme that is selectively activated in tissue macrophages and is markedly elevated in lysosomal storage diseases. A positive chitotriosidase is neither sensitive nor specific and can be useful in screening patients. It can be elevated in GD and NPD. CT levels above 200 nmol/h/ml are predictive for GD, NPD type A, B or C, but levels above 4000 nmol/h/ml are predictive of GD. (7,8) A diagnostic delay can occur in NPC due to overlapping

CONTACT Lavina Desai

Email : lavinadesai16@gmail.com

Address for Correspondence: Lavina Desai,
Seth G S Medical College, India.

©2018 Pediatric Oncall

features with Gaucher's disease and false positive testing for beta-glucosidase and hence requires a high index of suspicion along with mutation analysis to diagnose correctly.

Compliance with Ethical Standards

Funding: None

Conflict of Interest: None

References:

1. Nagral A. Gaucher's Disease. *J Clin Exp Hepatol.* 2014;4(1):37-50.
2. Mukherjee SM, Pandey M, Kapoor S, Padmapriya T. Infant with type A Niemann Pick Disease and Undetectable Niemann Pick cells in Bone Marrow. *Indian Pediatr.* 2012;49:490-492.
3. Gokulakrishnan V, Tamilselvan K. Niemann-Pick Disease available on <https://www.medindia.net/patientinfo/niemann-pick-disease.htm>. Reviewed on Oct 04,2017.
4. Vanier MT, Millat G. Niemann-Pick disease type C. *Clinical Genetics.* 2013;64:292.
5. Puri RD, Kapoor S, Krishnani PS, Dalal A, Gupta N, Muranjan M et al. Diagnosis and Management of Gaucher's Disease in India - Consensus guidelines of the Gaucher's Disease task force of the society for Indian Academy of Medical Genetics and the Indian Academy of Pediatrics. *Indian Pediatr.* 2018;55:143-153.
6. Mane V, Joshi RT. Niemann Pick Disease -A case report. *Journal of evolution of Medical and Dental Sciences.* 2012;1 (6).
7. Lo SM, McNamara J, Seashore MR, Mistry PK. Misdiagnosis of Niemann-Pick disease type C as Gaucher's disease. *Pediatrics.* 1977;(6):3-7.
8. Sheth JJ, Sheth FJ, Oza NJ, Gambhir PS, Dave UP, Shah RC. Plasma chitotriosidase activity in children with lysosomal storage disorders. *Indian J Pediatr.* 2010;77(2): 203-5.