

IMAGES IN CLINICAL PRACTICE

PRETERM WITH MILKY URINE

Anshika Taiwade¹, Shweta Anand¹, Rashmi Vishwakarma¹, Rajeev Sharma¹, G.C Bhatt².

¹Department of Pediatrics, LN Medical College and JK Hospital, Bhopal, Madhya Pradesh, India, ²Department Of Paediatric Nephrology, AIIMS, Bhopal, Madhya Pradesh, India.

KEYWORDS

Preterm, Milky Urine, Benign crystalluria.

A primigravida mother delivered a preterm (35±2 weeks) male baby by Emergency LSCS (Severe oligohydramnios) with birth weight of 2.28 kg. Baby required no resuscitation and shifted to mother side after routine care. Baby developed neonatal jaundice on 60 hours of life for which phototherapy started. Neonatal hyperbilirubinemia resolved within 24 hours and baby was discharged on 6th day of life. On 10th day mother came with complaint of passing turbid urine (Figure 1) and significant weight loss (300 gms).

Figure 1. Showing milky urine.



We hospitalized the baby and started initial management. Initial management of urine revealed amorphous amount of calcium oxalate crystal with no pus cell. septic screening were negative and urine culture showed commensal pathogen. Radioimaging done which was normal, RFT were normal. Urine calcium/creatinine ratio was high and urine chylomicron was negative.

Address for Correspondance: Anshika Taiwade, JK hospital, Kolar road, Bhopal, 462042, India.

Email: anshika22.taiwade@gmail.com

©2026 Pediatric Oncall

ARTICLE HISTORY

Received 19 January 2023

Accepted 27 July 2023

Figure 2. Breast feeding, Hydration improved.



Figure 3. Urine became clear



What is Benign crystalluria?

Finally we made diagnosis of benign crystalluria. Crystalluria refers to crystals found in urine when performing a urine test. Crystalluria is considered often as a benign condition and may represent Dehydration. Which in newborns can be corrected by encouraging breastfeeding.

Compliance with ethical standards

Funding: None

Conflict of Interest: None

References:

1. Milky urine in a premature infant by Lucia Marseglia & Sara Manti & Gabriella D'Angelo & Ignazio Barberi & Eloisa Gitto IPNA 2013.
2. Yamauchi S (1945) Chyluria: clinical, laboratory and statistical study of 45 personal cases observed in Hawaii. J Urol 54:318-347
3. Diamond E, Schapira HE (1985) Chyluria-a review of the literature. Urology 26:427-431
J Urol 54:318-347
3. Diamond E, Schapira HE (1985) Chyluria-a review of the literature. Urology 26:427-431
4. Hemal AK, Gupta NP (2002) Retroperitoneoscopic lymphatic management of intractable chyluria. J Urol 167:2473-2476
5. Koo CG, Van Langenberg A (1969) Chyluria. A clinical study. J R Coll Surg Edinb 14:31-41
6. WHO Expert Committee on Filariasis (1992) Lymphatic filariasis: the disease and its control. Fifth report of the WHO expert committee on Filariasis. WHO Tech Rep Ser 821:1-71
7. Buck AA (2002) Filariasis. In: Strickland GT (ed) Hunter's tropical medicine, 7th edn. WB Saunders, Philadelphia, pp 713-
8. Stalens JP, Falk M, Howmann-Giles R, Roy LP (1992) «Milky» urine-a child with chyluria. Eur J Pediatr 151:61-62
9. Kittredge RD, Hashim S, Roholt HB, Van Itallie TB, Finby N (1963) Demonstration of lymphatic abnormalities in a patient with chyluria. Am J Roentgenol 90:159-165
10. Punj J, Anand R, Darlong V, Pandey R (2013) Milky urine! A cause for concern? Indian J Anaesth 57:87-88

